DRAFT

12020 Olivine Basalt 312 grams



Figure 1: Photo of 12020. Scale in cm. NASA # S70-43639.

Introduction

12020 is an olivine basalt with large rounded olivine phenocrysts and elongate pyroxene crystals set in a variolitic groundmass of thin clinopyroxene and plagioclase laths (figure 2). It had one large zap pit on one surface.

Petrography

Klein et al. (1971) describe 12020 as a "mediumgrained olivine microgabbro consisting mainly of clinopyroxene, plagioclase and olivine. The clinopyroxene occurs as subhedral laths, up to several mm in length, as smaller anhderal grains, and as very thin, lathlike, crystals interleathed with plagioclase laths."

Lunar Sample Compendium C Meyer 2005



Figure 2: Photomicrograph of thin section 12020,13 illustrating round olivine and elongate pyroxene phenocrysts with variolitic intergrowths of plagioclase and pyroxen needles. NASA S70-30254. Field of view about 2 cm.





Mineralogy

Olivine: According to Klein et al. (1971) the cores of olivine in 12020 are rather homogeneous (Fo₇₀₋₇₇), whereas the rims range from Fo₇₀ to Fo₅₀ (figure 4). Kushiro et al. (1971) reported a wide range in olivine composition from Fo₇₄ to Fo₃.



Figure 4: Pyroxene and olivine composition of 12020 (adapted from Klein et al. 1971, Kushiro et al. 1971).

Pyroxene: Kushiro et al. (1971) and Klein et al. (1971) studied the composition of pyroxene in 12020 (figure 4). Pyroxene zones in Fe all the way to ferroheddenbergite (now there's a name).

Plagioclase: Plagioclase composition in 12020 range from An_{93} to An_{88} (Kushiro et al. 1971). Klein et al. (1971) report An_{98} to An_{80} , with the majority as An_{96} .

Chemistry

The rare earth element pattern is relatively flat (figure 5). 12020 is relatively Mg-rich (figure 6), apparently due to accumulation of olivine (Walker et al. 1976).

Radiogenic age dating

Alexander et al. (1972) reported an Ar/Ar age of 3.20 \pm 0.03 b.y. for 12020, consistent with that of other Apollo 12 basalts.

Mineralogica	l Mode for 1	2020	
-	Neal et	Klein et	Papike et
	al. 1994	al. 1971	al. 1976
Olivine	19	15.1	11.4
Pyroxene	51.2	58.6	61.4
Plagioclase	25.9	20	20.7
Opaques			5.6
Ilmenite	0.2		
Chromite +Usp	2.7	4.6	
Mesostasis	0.5	1.7	
"silica"			0.2

Cosmogenic isotopes and exposure ages

Hintenberger et al. (1971) determined exposure ages for 12020 using ³He (77 m.y.), ²¹Ne (71 m.y.) and ³⁸Ar (56 m.y.). The suntan age for 12020 (from etched solar flare track studies) is 2.6 m.y. (Bhandari et al. 1971).

Other Studies

Bogard et al. (1971) reported the content and isotopic composition of rare gases in 12020.

List of Photo #s for 12020

S69-24225	TS
S69-24213	closeup
S69-64130	color mug
S69-64105	
\$69-63324-332	B&W mug
S70-43638-640	color mug
S70-19641-644	wire saw cut
S70-49135-144	TS color
S70-25406-408	
S70-25421-424	
S70-30251-253	
S70-25890-893	
S70-27991	
S70-31559-566	TS



Figure 5: Rare-earth-element diagram for 12020 (idms data from Hubbard connected).



Figure 6: Composition of lunar basalts showing 12020.



Table 1. Chemical composition of 12020.

reference weight	Kushiro71		Hubbard71		Weisman	n75	75 Wanke71		Cuttitta71		Wakita	a71		Haskin	71	Compston7		Anders	71
SiO2 % TiO2 Al2O3 FeO MnO MgO CaO Na2O K2O P2O5 S % sum	44.45 2.54 7.99 20.65 0.26 14.89 8.53 0.21 0.06 0.02	(a) (a) (a) (a) (a) (a) (a) (a) (a)	0.19 0.056	(b)	0.056	(b)	43.86 2.64 7.2 21.1 0.28 15.65 8.12 0.17 0.046	(c) (c) (c) (c) (c) (c) (c) (c)	44.6 2.56 8 20.7 0.27 14.4 8.53 0.23 0.06 0.08	(d) (d) (d) (d) (d) (d) (d) (d) (d)	42.2 2.7 8.5 21.8 0.253 16.1 8.7 0.22	3 8.3 0.26 8.8 0.213 0.069	(c) (c) (c) (c) (c) (c) (c) (c)			44.66 2.73 7.31 21.58 0.28 13.91 8.73 0.21 0.064 0.08 0.06	(f) (f) (f) (f) (f) (f) (f) (f) (f) (f)		
Sc ppm V Cr Co Ni Cu Zn Ga Ge ppb As Se Rb Sr Y Zr Nb Mo Ru Rh	4653	(a)					45.4 4560 61 6.9	(c) (c) (c) (c)	39 155 4330 64 77 9 4.8	(d) (d) (d) (d) (d) (d) (d)	42 180 4187 61	200	(c) (e) (c) (c)			146 3780 50 50 13 4 1.8	(f) (f) (f) (f) (f) (f)	68 0.74	(e) (e)
			0.997 93.6	(b) (b)	0.997 93.6	(b) (b)			1.4 65 37 119 13	(d) (d) (d) (d)		1 34	(e) (e)			1.03 91.4 32 97 5	(f) (f) (f) (f) (f)	0.114 0.85	(e) (e)
Pd ppb Ag ppb Cd ppb In ppb Sn ppb Sb ppb												12	(e)					0.98 1.1 2.7	(e) (e) (e)
le ppb Cs ppm Ba La			64.4	(b)	64.4	(b)	4 82	61 ((c)	(d)	25 5.9	0.05 (e (c 5.6 (e	(e) (c) (e)	 ⇒) ⇒) ⇒) 5.19 (c) 		60) 4	(f) (f)	0.039	(e)	
Ce			16.1	(b)	16.1	(b)					0.0	16.1	(e) (e)	14.5	14.5 (c)	ý 11	(f)		
Pr Nd Sm Eu Gd Tb Dy Ho Er			12 4.5 0.839 5.43 6.13	(b) (b) (b) (b) (b)	12 4.5 0.839 5.43 6.13	(b) (b) (b) (b) (b)	3.4 0.82 0.91 5.68 1.07	(c) (c) (c) (c) (c)			4.08 0.79	2.2 12 4.1 0.76 5.4 0.96 5.2 1.55 3.5	 (e) 	13 3.92 0.76 5.3 1.02 7.1 1.34 3.8	(c) (c) (c) (c) (c) (c) (c)				
Tm			2.60	(b)	2 20	(b)	2.01	(0)	5 1	(d)	20	0.59	(e) (e)	2 4 2	(0)				
Yb Lu Hf Ta W ppb Re ppb			3.69	(U) 60	0.14	(b) (b)	0.42 3.8 0.45	(c) (c) (c) (c)	J. I	(u)	3.8 0.54 2.4	3.7 0.52	(e) (e) (c)	3.43 0.51	3.43 (c)).51 (c))			
lr ppb																		0.04	(e)
Pt ppb Au ppb													(c)					0.36	(e)
0 ppm technique	(a) con	venti	ional we	t, (b)) IDMS, (c) IN	AA , (d)	mixe	ed mciro	cher	n, XRF,	emissioi	n spe	ec., (e) R	NAA	A, (f) XRF			



Lunar Sample Compendium C Meyer 2005